=> d his (FILE 'HOME' ENTERED AT 14:45:59 ON 15 JUL 2003) FILE 'MEDLINE, AGRICOLA, CANCERLIT, SCISEARCH, CAPLUS, MEDICONF' ENTERED AT 14:46:22 ON 15 JUL 2003 L11852 S TRANSGENIC (L) (CHICKEN OR HEN OR FOWL OR BIRD) 72 S L1 AND (INTERFERON OR ERYTHROPOIETIN OR EPO OR IFN) L2 39 DUP REM L2 (33 DUPLICATES REMOVED) L3 17 S L3 AND PY<=1997 L4 17 SORT L4 PY 1.5 FILE 'STNGUIDE' ENTERED AT 14:52:25 ON 15 JUL 2003 FILE 'MEDLINE, AGRICOLA, CANCERLIT, SCISEARCH, CAPLUS, MEDICONF' ENTERED AT 14:54:01 ON 15 JUL 2003 E IVARIE ROBERT?/AU 21 S E2 3 S L6 AND TRANSGEN? L7 L8 3 DUP REM L7 (0 DUPLICATES REMOVED) L9 6 S L6 AND TRANS? 1.10 29 S E1 17 S L10 AND TRANS? L11 5 S L11 AND EGG? L12=> d an ti so au ab pi 112 1-5 L12 ANSWER 1 OF 5 2002719100 MEDLINE AN TI Avian transgenesis: progress towards the promise. TRENDS IN BIOTECHNOLOGY, (2003 Jan) 21 (1) 14-9. Ref: 76 Journal code: 8310903. ISSN: 0167-7799. ΑU Ivarie Robert The hen has long held promise as a low cost, high-yield bioreactor for the AB production of human biopharmaceuticals in egg whites. A typical egg white contains 3.5-4.0 grams of protein, more than half of which comes from a single gene (ovalbumin). Harnessing the power of the gene to express a recombinant protein could yield up to a gram or more of the protein in the naturally sterile egg. Accordingly, a major effort has been underway for more than a decade to develop robust methods for modification of the chicken genome. This effort intensified in the mid-1990s when several avian transgenic companies entered the scene. Progress has been made in that time but much remains to be done. L12 ANSWER 2 OF 5 MEDLINE MEDLINE ΔN 2002190519 Expression of exogenous protein in the egg white of transgenic chickens. SO NATURE BIOTECHNOLOGY, (2002 Apr) 20 (4) 396-9. Journal code: 9604648. ISSN: 1087-0156. Harvey Alex J; Speksnijder Gordon; Baugh Larry R; Morris Julie A; AU Ivarie Robert AR Using a replication-deficient retroviral vector based on the avian leukosis virus (ALV), we inserted into the chicken genome a transgene encoding a secreted protein, beta-lactamase, under the control of the ubiquitous cytomegalovirus (CMV) promoter. Biologically active beta-lactamase was secreted into the serum and egg white of four generations of transgenic chickens. The expression levels were similar in successive generations, and expression levels in the magnum of the oviduct were constant over at least 16 months in transgenic hens, indicating that the transgene was stable and not subject to silencing. These results support the potential of the hen as a bioreactor for the production of commercially valuable, biologically active proteins in egg white. L12 ANSWER 3 OF 5 CAPLUS COPYRIGHT 2003 ACS 2002:931060 CAPLUS AN DN 138:215789 ΤI Avian transgenesis: progress towards the promise Trends in Biotechnology (2002), Volume Date 2003, 21(1), 14-19 SO

CODEN: TRBIDM; ISSN: 0167-7799

AU Ivarie, Robert

AB A review. The hen has long held promise as a low cost, high-yield bioreactor for the prodn. of human biopharmaceuticals in egg whites. A typical egg white contains 3.5-4.0 g of protein, more than half of which comes from a single gene (ovalbumin). Harnessing the power of the gene to express a recombinant protein could yield up to a gram or more of the protein in the naturally sterile egg.

Accordingly, a major effort has been underway for more than a decade to develop robust methods for modification of the chicken genome. This effort intensified in the mid-1990s when several avian transgenic companies entered the scene. Progress has been made in that time but much remains to be done.

L12 ANSWER 4 OF 5 CAPLUS COPYRIGHT 2003 ACS

AN 2002:291304 CAPLUS

DN 137:135738

TI Expression of exogenous protein in the egg white of transgenic chickens

SO Nature Biotechnology (2002), 20(4), 396-399 CODEN: NABIF9; ISSN: 1087-0156

AU Harvey, Alex J.; Speksnijder, Gordon; Baugh, Larry R.; Morris, Julie A.; Ivarie, Robert

AB Using a replication-deficient retroviral vector based on the avian leukosis virus (ALV), the authors inserted into the chicken genome a transgene encoding a secreted protein, .beta.-lactamase, under the control of the ubiquitous cytomegalovirus (CMV) promoter. Biol. active .beta.-lactamase was secreted into the serum and egg white of four generations of transgenic chickens. The expression levels were similar in successive generations, and expression levels in the magnum of the oviduct were const. over at least 16 mo in transgenic hens, indicating that the transgene was stable and not subject to silencing. These results support the potential of the hen as a bioreactor for the prodn. of com. valuable, biol. active proteins in egg white.

L12 ANSWER 5 OF 5 CAPLUS COPYRIGHT 2003 ACS

AN 2000:145005 CAPLUS

DN 132:204036

TI Direct avian oviduct **transgenesis** for exogenous protein expression in poultry **eggs**

SO PCT Int. Appl., 54 pp. CODEN: PIXXD2

IN Ivarie, Robert; Harvey, Alex J.; Murphy, George F., Jr.; Rapp, Jeffrey C.

AB Methods for prepg. transgenic avians which express exogenous protein substantially only in their oviducts are disclosed. Each of the methods comprises delivering nucleic acid expression cassettes directly to the oviducts of the avians. The exogenous protein expressed by the expression cassette is secreted into the lumen of the avian oviduct and secreted into the eggs of the transgenic avians.

Methods for prepg. eggs which contain exogenous protein, such as human interferon, and methods for the prodn. of proteins are also disclosed. The methods for direct oviduct transgenesis may also be used to assess the suitability of expression cassettes or exogenous proteins for expression in the avian oviduct.

APPLICATION NO. DATE PATENT NO. KIND DATE ----PΙ WO 2000011151 20000302 WO 1999-US19393 19990825 A2 А3 WO 2000011151 20000615 W: AE, AL, AM, AT, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, CZ, DE, DE, DK, DK, DM, EE, EE, ES, FI, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM RW: GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG AU 9956907 A1 20000314 AU 1999-56907 19990825

(FILE 'HOME' ENTERED AT 14:45:59 ON 15 JUL 2003) FILE 'MEDLINE, AGRICOLA, CANCERLIT, SCISEARCH, CAPLUS, MEDICONF' ENTERED AT 14:46:22 ON 15 JUL 2003 L11852 S TRANSGENIC (L) (CHICKEN OR HEN OR FOWL OR BIRD) 72 S L1 AND (INTERFERON OR ERYTHROPOIETIN OR EPO OR IFN) 1.2 L3 39 DUP REM L2 (33 DUPLICATES REMOVED) L417 S L3 AND PY<=1997 L5 17 SORT L4 PY FILE 'STNGUIDE' ENTERED AT 14:52:25 ON 15 JUL 2003 FILE 'MEDLINE, AGRICOLA, CANCERLIT, SCISEARCH, CAPLUS, MEDICONF' ENTERED AT 14:54:01 ON 15 JUL 2003 => d an ti so au ab pi 15 8 ANSWER 8 OF 17 CAPLUS COPYRIGHT 2003 ACS AN 1995:645229 CAPLUS DN 123:49258 TT Chicken interferon gene, its sequence and recombinant expression, and its antiviral action SO PCT Int. Appl., 46 pp. CODEN: PIXXD2 IN Sekellick, Margaret J.; Marcus, Philip I.; Ferrandino, Anthony F. AB A chicken interferon gene sequence and its deduced amino acid sequence is provided. There is also disclosed a method of producing biol. active chicken interferon recombinantly, a method of isolating other non-mammalian interferon genes, a method of making a transgenic fowl having the chicken interferon gene incorporated therein, and a method for delivery of the chicken interferon in the bird, such as by genetic immunization or aerosol. Expression of the gene yields functional chicken interferon. Thus, chicken interferon cDNA was isolated which coded for the complete mature interferon sequence of 162 amino acids plus a 31-residue signal peptide. A cDNA probe was also isolated, consisting of 269 nucleotides with a 5'-primer region of 32 bases, a 3' primer region of 20 bases, and a 217-base partial sequence of the chicken interferon gene. The probe can be used in screening a non-mammalian cDNA library by std. procedures. Glycosylated recombinant chicken interferon could be expressed in transfected mouse L(Y) cells, human WISH cells, and COS-1 cells, and displayed antiviral activity. Chicken interferon cDNA can be integrated into plasmid or retroviral vectors under the control of the **chicken** metallothionein promoter (for Cd or Zn induction) and introduced into sperm, egg, or zygote of fowls in order to prevent viral and/or parasitic infection. PATENT NO. KIND DATE APPLICATION NO. DATE -----WO 9511302 PΤ A1 19950427 WO 1994-US11660 19941021 <--W: AU, BR, CA, JP, KR RW: AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE

19970624

19990323

19950508

19980108

19960807

R: AT, BE, CH, DE, DK, ES, FR, GB, IT, LI, NL

Α

Α

A1

B2

A1

US 5641656

US 5885567

AU 9480771

EP 724638

AU 684849

SK-1636

US 1993-139909 19931022 <--

19940919

19941021 <--

19941021 <--

US 1994-308821

AU 1994-80771

EP 1994-931841

=> d his

AU 752946

EP 1023442

JP 2001520009

IE, FI

```
(FILE 'HOME' ENTERED AT 14:45:59 ON 15 JUL 2003)
      FILE 'MEDLINE, AGRICOLA, CANCERLIT, SCISEARCH, CAPLUS, MEDICONF' ENTERED
     AT 14:46:22 ON 15 JUL 2003
            1852 S TRANSGENIC (L) (CHICKEN OR HEN OR FOWL OR BIRD)
L1
              72 S L1 AND (INTERFERON OR ERYTHROPOIETIN OR EPO OR IFN)
L2
              39 DUP REM L2 (33 DUPLICATES REMOVED)
L3
              17 S L3 AND PY<=1997
L4
              17 SORT L4 PY
L5
     FILE 'STNGUIDE' ENTERED AT 14:52:25 ON 15 JUL 2003
      FILE 'MEDLINE, AGRICOLA, CANCERLIT, SCISEARCH, CAPLUS, MEDICONF' ENTERED
     AT 14:54:01 ON 15 JUL 2003
                  E IVARIE ROBERT?/AU
              21 S E2
L6
L7
                3 S L6 AND TRANSGEN?
L8
                3 DUP REM L7 (0 DUPLICATES REMOVED)
=> d an ti so au ab pi 18 2
     ANSWER 2 OF 3 CAPLUS COPYRIGHT 2003 ACS
L8
AN
     1999:271485 CAPLUS
DN
     130:277660
     Vectors comprising a magnum-specific promoter for avian
      transgenesis
SO
     PCT Int. Appl., 67 pp.
     CODEN: PIXXD2
     Ivarie, Robert D.; Harvey, Alex J.; Morris, Julie A.; Liu,
TN
     This invention provides vectors and methods for the stable introduction of
AB
      exogenous nucleic acid sequences into the genome of a bird and for
      expressing said exogenous sequences to alter the phenotype of the bird or
      to produce desired proteins. In particular, transgenic chickens
      are produced which express exogenous sequences in their oviducts.
      which contain exogenous proteins are also produced. In one specific
      embodiment, an avian leukosis virus retroviral vector is used which
      comprises a modified pNLB plasmid contg. an exogenous gene that is
      inserted downstream of a segment of the ovalbumin promoter region.
      total length of the ovalbumin promoter segment may be from about 0.88 kb
      to about 7.4 kb in length, and includes both the steroid-dependent
      regulatory element and the neg. regulatory element. An RNA copy of the
      modified retroviral vector, packaged into viral particles is used to
      infect embryonic blastoderms which develop into transgenic
      birds. Alternatively, helper cells which produce the retroviral
      transducing particles are delivered to the embryonic blastoderm.
      PATENT NO.
                        KIND DATE
                                                APPLICATION NO. DATE
                        ----
     WO 9919472
                         A1 19990422
                                                WO 1998-US21975 19981015
PΙ
          W: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE,
              DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW,
              MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR,
              \mathtt{TT},\ \mathtt{UA},\ \mathtt{UG},\ \mathtt{UZ},\ \mathtt{VN},\ \mathtt{YU},\ \mathtt{ZW},\ \mathtt{AM},\ \mathtt{AZ},\ \mathtt{BY},\ \mathtt{KG},\ \mathtt{KZ},\ \mathtt{MD},\ \mathtt{RU},\ \mathtt{TJ},\ \mathtt{TM}
          RW: GH, GM, KE, LS, MW, SD, SZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG
      CA 2307840
                         AA 19990422
                                                CA 1998-2307840 19981015
     AU 9911899
                          Α1
                               19990503
                                                AU 1999-11899
                                                                    19981015
```

R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT,

EP 1998-954994

JP 2000-516025

19981015

19981015

20021003

20000802

20011030

B2

A1

T2

L Number	Hits	Search Text	DB	Time stamp
1	1695	Transgenic WITH (chicken OR bird or fowl	USPAT;	2003/07/15 14:20
		or turkey or hen)	US-PGPUB; EPO; JPO; DERWENT	
7	661	(Transgenic WITH (chicken OR bird or fowl or turkey or hen)) and egg	USPAT; US-PGPUB; EPO; JPO; DERWENT	2003/07/15 14:12
13	16	((Transgenic WITH (chicken OR bird or fowl or turkey or hen)) and egg) and (egg SAME interferon)	USPAT; US-PGPUB; EPO; JPO; DERWENT	2003/07/15 14:13
19	7	((Transgenic WITH (chicken OR bird or fowl or turkey or hen)) and egg) and (egg SAME erythropoietin)	USPAT; US-PGPUB; EPO; JPO; DERWENT	2003/07/15 14:13
25	. 21	(((Transgenic WITH (chicken OR bird or fowl or turkey or hen)) and egg) and (egg SAME interferon)) or (((Transgenic WITH (chicken OR bird or fowl or turkey or hen)) and egg) and (egg SAME erythropoietin))	USPAT; US-PGPUB; EPO; JPO; DERWENT	2003/07/15 14:13
31	28		USPAT; US-PGPUB; EPO; JPO; DERWENT	2003/07/15 14:20
38	16	(US-6395961-\$ or US-6333192-\$ or US-6156569-\$ or US-5897998-\$ or US-5784992-\$ or US-5162215-\$ or US-H001065-\$ or US-4997763-\$ or US-6515199-\$).did. or (US-20020116732-\$ or US-20020108132-\$ or US-20020028488-\$ or US-20010039668-\$ or US-20030126629-\$ or US-20030126628-\$ or US-20030074681-\$).did.	USPAT; US-PGPUB	2003/07/15 14:29
41	16	((US-6395961-\$ or US-6333192-\$ or US-6156569-\$ or US-5897998-\$ or US-5784992-\$ or US-5162215-\$ or US-H001065-\$ or US-4997763-\$ or US-6515199-\$).did. or (US-20020116732-\$ or US-20020108132-\$ or US-20020028488-\$ or US-20010039668-\$ or US-20030126629-\$ or US-20030074681-\$).did.) and stage	USPAT; US-PGPUB; EPO; JPO; DERWENT	2003/07/15 14:29
-	4905	800/\$?.ccls.	USPAT; US-PGPUB; EPO; JPO; DERWENT	2002/09/16 13:07
-	835	800/\$?.ccls. and (chick\$10 or bird or fowl)	USPAT; US-PGPUB; EPO; JPO; DERWENT	2002/09/16 13:25
-	34	(800/\$?.ccls. and (chick\$10 or bird or fowl)) and (transgen\$10 ADJ (chick\$10 or bird or fowl))	USPAT; US-PGPUB; EPO; JPO; DERWENT	2002/09/16 13:08
-	12	(US-4997763-\$ or US-H001065-\$ or US-5162215-\$ or US-5897998-\$ or US-6156569-\$ or US-6333192-\$ or US-6395961-\$ or US-5784992-\$).did. or (US-20010039668-\$ or US-20020028488-\$ or US-20020108132-\$ or US-20020116732-\$).did.	USPAT; US-PGPUB	2002/09/16 13:21
-	2	((US-4997763-\$ or US-H001065-\$ or US-5162215-\$ or US-5897998-\$ or US-6156569-\$ or US-6333192-\$ or US-6395961-\$ or US-5784992-\$).did. or (US-20010039668-\$ or US-20020028488-\$ or US-20020108132-\$ or US-20020116732-\$).did.) and (EFO or interferon\$10)	USPAT; US-PGPUB; EPO; JPO; DERWENT	2002/09/16 13:25

[-	51	(800/\$?.ccls. and (chick\$10 or bird or	USPAT;	2002/09/16 13:27
		fowl)) and EPO	US-PGPUB;	
	i		EPO; JPO;	
			DERWENT	1
-	1213	EPO and (chick\$10 or bird or fowl)	USPAT;	2002/09/16 13:26
}			US-PGPUB;	
			EPO; JPO;	
1_	486	(FDO and (abiabata and a second	DERWENT	
-	400	(ter term (term term term term term term term term	USPAT;	2002/09/16 13:26
		transgenic\$10	US-PGPUB;	!
			EPO; JPO;	
_	31	(800/\$?.ccls. and (chick\$10 or bird or	DERWENT	0000/00/45 40 01
		fowl)) and Erythropoietin\$5	USPAT; US-PGPUB;	2002/09/16 13:34
		1 - 0.12// and bry enroporeering	EPO; JPO;	
	•		DERWENT	
-	110	(800/\$?.ccls. and (chick\$10 or bird or	USPAT:	2002/09/16 13:42
		fowl)) and interferon\$5	US-PGPUB;	2002/03/16 13:42
		•	EPO; JPO;	
			DERWENT	
-	300	t tree=== t and tonical of Dild of	USPAT;	2002/09/16 13:42
		fowl)) and egg\$2	US-PGPUB;	3, 13, 13 13, 12
			EPO; JPO;	
1	e =		DERWENT	
-	65	((800/\$?.ccls. and (chick\$10 or bird or	USPAT;	2002/09/16 13:43
		fowl)) and egg\$2) and (interferon or	US-PGPUB;	
		erythropoietin)	EPO; JPO;	
L			DERWENT	